

UNIVERSITY OF TORONTO LIBRARIES

A standard linear barcode consisting of vertical black lines of varying widths on a white background.

3 1761 01291693 8

Digitized by the Internet Archive
in 2008 with funding from
Microsoft Corporation

23

2

224?

ILLUSTRATIONS
OF
BRITISH FUNGI.



ot
halloph
C

ILLUSTRATIONS

OF

BRITISH FUNGI

(HYMENOMYCETES),

To serve as an Atlas to the Handbook of "British Fungi."

BY

M. C. COOKE, M.A., LL.D., A.L.S.,

Late President of the Quekett Microscopical Club. President of the Hackney Natural History and Microscopical Society. Fellow of the Botanical Society of Edinburgh. Corresponding Member of the Academy of Natural Sciences, Philadelphia; of the Academy of Science of New York; of the Portland Society of Natural History, Maine, and of the Società Crittogramologica Italiana. Honorary Member of the Cryptogamic Society of Scotland; the Woolhope Field Club; the Hertfordshire Natural History Society; the Norfolk and Norwich Naturalists' Society; the Essex Field Club; and other Societies.

VOL. VI.

London:

WILLIAMS AND NORGATE, HENRIETTA STREET, W.C.

1886-1888.


131426
61111


QK

607

C67

V.6










M.C.C.

CORTINARIUS (DERMOCYBE) OCHROLEUCUS. Fries.
on the ground. Epping Forest.












CORTINARIUS (DERMOCYBE) DECUMBENS. Fries.
grassy places in woods. Epping Forest.



CORTINARIUS (DERMOCYBE) DIABOLICUS. Fries.
under beech trees. Dorking. Nov. 1880.








CORTINARIUS (DERMOCYBE) TABULARIS. Bull.
in mixed woods, near Carlisle. 1886.











M.C.C.

CORTINARIUS (DERMOCYBE) CAMURUS. Fries.
on ground in Epping Forest, Chingford. Nov. 1882.









MCC.


CORTINARIUS (*DERMOCYBE*) CANINUS. Fries.
on the ground. Tunnel Woods, Watford. 1883.






CORTINARIUS (DERMOCYBE) MYRTILLINUS. Fries.
under beech trees. Christchurch, Hants.









CORTINARIUS (INOLOMA) AZUREUS. Fries.
Amongst leaves. Alresford, Hants.











M.C.C.

CORTINARIUS (DERMOCYBE) ALBO-CYANEUS. Fries.
under trees. Pontrilas. Sept. 1885.










M.J.B.

CORTINARIUS (DERMOCYBE) ANOMALUS. Fries.
on the ground in woods. Lyndhurst, Haywood Forest, &c.











CORTINARIUS (DERMOCYBE) LEPIDOPUS. Cooke.
in healthy places; Epping Forest, and near Carlisle. Sept. 1887.









A




CORTINARIUS (DERMOCYBE) MILTINUS. Fries.
amongst moss. Ledbury. Oct. 1883.


B




odour raphanoid.




CORTINARIUS (DERMOCYBE) CINNABARINUS. Fries.
in beech woods. near Hereford.





CORTINARIUS (DERMOCYBE) SANGUINEUS. Fries.
on the ground in Epping Forest. Nov. 1882.






CORTINARIUS (DERMOCYBE) ANTHRACINUS. Fries.
in woods. Coed Coch.




CORTINARIUS (DERMOCYBE) ORELLANUS Fries.
in woods. Coed Coch.






M.C.C.


CORTINARIUS (DERMOCYBE) CINNAMOMEUS. Fries,
on the ground in woods. Dimore.





M.C.C.

CORTINARIUS (DERMOCYBE) CINNAMOMEUS. Fries, variety.
on the ground. near Carlisle. 1886.






M.C.C.

CORTINARIUS (DERMOCYBE) CINNAMOMEUS. Fries.

var. β. SEMI-SANGUINEUS. Fries.


on the ground. Lyme Regis. Oct. 1883.






A

CORTINARIUS (DERMOCYBE) CINNAMOMEUS. Fries.
var. CROCEUS. Fr. Haywood Forest. 1881.



B


CORTINARIUS (DERMOCYBE) CROCEO-CONUS. Fries.
in pine woods. Wyelliffe wood.





M.J.B.


CORTINARIUS (DERMOCYBE) ULIGINOSUS. — Berk.
in boggy woods. — King's Cliffe.





CORTINARIUS (*DERMOCYBE*) INFUCATUS
on the ground in woods. — Haywood Forest


PL. 781.
Aug. 1883.





Boudier


CORTINARIUS (DERMOCYBE) COTONEUS Fries
under oaks, or in mixed woods.



odour strong.



CORTINARIUS (DERMOCYBE) SUBNOTATUS. Pers.
amongst fir leaves. Alresford, Hants. Nov. 1886.





C.B.

CORTINARIUS (*DERMOCYBE*) VALGUS. Fries. (var.)
in woods near Bristol. Oct. 1856.



odour strong




CORTINARIUS (DERMOCYBE) RAPHANOIDES. Fries.
in mixed woods. Fordingbridge.

B



CORTINARIUS (DERMOCYBE) VENETUS. Fries.
under beech trees. Aboyne.





C.B.


CORTINARIUS (TELAMONIA) MACROPUS. Fries.
on the ground, near Bristol. 1884.





C.B.

CORTINARIUS (*ELAMIA*) LANIGER Fries.
under trees. — *Sa du Linn.* — *Bristol.* — Sept. 1885.






strong scented.




CORTINARIUS (TELAMONIA) BIVELUS Fries.
in woods. Tenedor. Oct. 1881.






CORTINARIUS (*TELAMONIA*) BULBOSUS. Sow.
in woods. Wycliffe Wood. Oct. 1883.






CORTINARIUS (TELAMONIA) URBICULUS. Fries.
amongst grass. Cabalva. Oct. 1879.





C.B.

CORTINARIUS (TELAMONIA) LICINIPES. Fries. var. *ROBUSTIOR.*
in damp woods. Blaise Castle. Oct. 1884.






M.C.C.

CORTINARIUS (*TELAMONIA*) MICROCYCLUS. Fries.
under trees. Broxbourne. Oct 1887.

has been allowed to be the
same species as that in
pl. 801. It suits H. &
t 157 f 2 but not Green
Ridge t 370. I have
very great confidence in
the plate".




CORTINARIUS (TELAMONIA) TORVUS — Fris.
on the ground in woods. — Wych-Tree Wood, Oct. 1882






CORTINARIUS (TELAMONIA) IMPENNIS. Fries.
under fir trees. Dinedor. Oct. 1881.



odour strong.




CORTINARIUS (TELAMONIA) SCUTULATUS.
under trees, Wycliffe Wood. Fr.

CORTINARIUS (TELAMONIA) QUADRICOLOR. Fr.
under beech. Blaise Castle, Bristol. Oct. 1882.




CORTINARIUS (TELAMONIA) EVERNIUS. Fries. (Type).
in damp places. Epping Forest, 1855.






M.C.C.

CORTINARIUS (TELAMONIA) QUADRICOLOR, Frise
amongst grass. near Brusonene. Oct. 1887






CORTINARIUS (*TELAMONIA*) ARMILLATUS. Fries.
in mixed woods. Wycliffe Wood, Ludlow.






CORTINARIUS (TELAMONIA) HÆMATOCHELIS. Bull.
in woods. Coal Coch, Abergavenny.






CORTINARIUS (TELAMONIA) LIMONIUS. Fries.
on ground in woods. near Bristol.




CORTINARIUS (TELAMONIA) HELVOLUS. Bull
in woods. Durham Downs. Oct. 1883.






CORTINARIUS (TELAMONIA) HINULEUS. — Fries.
on the ground, in woods. — Dior. Oct. 1881.






CORTINARIUS (TELAMONIA) GENTILIS. Fries.
on the ground. Haywood Forest, and Rothiemurchas.





M.C.C.


CORTINARIUS (TELAMONIA) HELVELLOIDES. Fries.
amongst beech leaves. Christchurch.



flesh spongy.




MARASMIUS LEPTOCEPHALUS Vahl.
var. *leptocephalus*. - *Leptocephalus*.




odour nitrous.


M.C.C.

CORTINARIUS (TELAMONIA) NITROSUM. Cooke.
on the ground. Durdham Down, Bristol.






CORTINARIUS (TELAMONIA) INJUCUNDUS. Weinm.
under trees. Swarraton, Hants. Nov. 1886.





CORTINARIUS (TELAMONIA) BRUNNEUS. Fries.
under trees. Hollywood Forest, 1882.






M.C.C.


CORTINARIUS (TELAMONIA) BRUNNEUS
on the ground. Epping Forest.

Fri. (Tigr.)
Oct. 29, 1887






CORTINARIUS (TELAMONIA) GLANDICOLOR. *Fries.*
amongst moss. Fordingbridge. 1883.






CORTINARIUS (TELAMONIA) PUNCTATUS. Fries.
under beech. near Ascot, 1881.



hygrophanous.

CORTINARIUS (TELAMONIA) TRIFORMIS var. SCHAEFFERI. Fries.
on the ground. Alresford, Hants Oct. 1885.







forma exannulatus.

12-18 X 6

CORTINARIUS (TELAMONIA) BIFORMIS. Fries.
in mixed woods. Park End, Forest of Dean. Oct. 1887.





CORTINARIUS (TELAMONIA) PERISCELIS. Weinm.
in boggy places. Near Bristol, &c.




CORTINARIUS (TELAMONIA) FLEXIPES. Fries.
in fir woods. Dimmore.

odour strong.



CORTINARIUS (TELAMONIA) FLABELLUS. Fries.
under beech. Codd Coch.





CORTINARIUS (TELAMONIA) PSAMMOCEPHALUS. Bull.
under firs. Credenhill Camp, 1882.




CORTINARIUS (TELAMONIA) ILEOPODIUS. Bull.
in beech woods. near Bristol.



M.C.C.

CORTINARIUS TELAMENEUS INCISUS. — Fr. — var. B
On the soil. — Presently fruiting. K.W.





CORTINARIUS (TELAMONIA) HEMITRICHUS. *Fruxt.*
amongst moss, &c. Belmont, Hereford, 1886.




strong scented.

CORTINARIUS (TELAMONIA) RIGIDUS. Fries.
in damp woods. near Carlisle. 1886.



M.C.C.


CORTINARIUS (TELAMONIA) PALEACEUS Fries.
in moist woods. Haywood Forest. Oct. 1854.






C.B.

CORTINARIUS (HYDROCYBE) FIRMUS. *Iris.*
on the ground. near Bristol.






CORTINARIUS (HYDROCYBE) SUBFERRUGINEUS. — Fries.
on the ground in woods. — Haywood Forest, Sept. 1853






CORTINARIUS (HYDROCYBE) ARMENIACUS. Schaff.
in fir woods. Mattishall. Aug. 1853.



*often cæspitose.**aerid.**variety.*


CORTINARIUS (HYDROCYBE) DAMASCENUS. Fries.
under trees. Pontrilas. Sept. 1885.





M.C.C.


CORTINARIUS (HYDROCYBE) PRIVIGNUS. — *Fres.*
under fir trees. — *Diamond.* — 1885.





CORTINARIUS (HYDROCYBE) DURACINUS. Fries.


in woods. Hereford Fungus Show, 1881.





G.M.


CORTINARIUS (HYDROCYBE) ILLUMINUS. Fries.
under firs. near Scarborough.



stained with red.

G.M.


COPTODISCUS (R. RETICULARIS) RETICULATUS Dofle
in Linnemann's Archiv für Naturforschung





CORTINARIUS (HYDROCYBE) DILUTUS.
on the ground in Epping Forest, Chingford.

Fries.
Nov. 1882.




odour faint.



M.C.C.


CORTINARIUS (HYDROCYBE) SATURNINUS. Fries.
amongst grass. Donaore 1886.



subcaspitose.


CORTINARIUS (HYDROCYBE) IMBUTUS. Fries.

in mixed woods. Forres, N.B. Sept. 1887






CORTINARIUS (HYDROCYBE) CASTANEUS. Fries.
under trees. Kew Gardens, and Hereford.





CORTINARIUS (*HYDROCYBE*) BICOLOR. Cooke.
in mixed woods. Park End, Forest of Dean. Oct. 1887.






8X 4-5


M.C.C.

CORTINARIUS (HYDROCYBE) BALAUSTINUS. Fries.
under beech. Credinhill Camp. Oct. 1882






CORTINARIUS (HYDROCYBE) COLUS. Fr.
on the ground. Blaize Castle, and Whittington.





CORTINARIUS (HYDROCYBE) ISABELLINUS. Fries.
in pine woods Ascot.





M.C.C.


cœspitose.

CORTINARIUS (HYDROCYBE) RENIDENS. Pries.
under trees. Highbeach, Epping Sept. 1882.






CORTINARIUS (*HYDROCYBE*) URACEUS. Fries.
under fir trees, near Hereford.





M.C.C.

CORTINARIUS (HYDROCYBE) JUBARINUS. Fries.
amongst grass. Lyndhurst, Hants. Oct. 1883.




pileus shining.



M.C.C.

CORTINARIUS (HYDROCYBE) PATERIFORMIS. var MAJOR. Fries.
amongst grass Neasden, Middlesex. Sept. 1881




pileus shining.

M.C.C.

CORTINARIUS (HYDROCYBE) UNIMODUS. *Britzelmeyer.*
in grassy places. Carlisle. Sept. 1887.




CORTINARIUS (HYDROCYBE) DOLABRATUS. Fries.
on the ground amongst firs. Dereham, Norfolk. Nov. 1853.






CORTINARIUS (HYDROCYBE) RIGENS. Fries.
under fir. Epting Forest. June, 1869





G.M.

CORTINARIUS (HYDROCYBE) KROMBHLZII. Fries.
amongst moss. Scarborough, Yorks.







Hussey.

CORTINARIUS (HYDROCYBE) REEDII. Berk.
amongst moss. Hayes Common. Spring.

B




CORTINARIUS (HYDROCYBE) LEUCOPUS. Bull.
under fir trees. near Kendal.





G.M.

CORTINARIUS (HYDROCYBE) SCANDENS. Fr. Mon. (nec. Fr. Icon.)
on moist ground. near Downham, Norfolk.



A



CORTINARIUS (HYDROCYBE) ERYTHRINUS. Fries.
amongst leaves. Wycliffe Wood, Oct. 1883.

B



CORTINARIUS (HYDROCYBE) DECIPIENS. Fries.
on the ground, in woods. Norfolk. 1852.


strong odour.



hygrophanous.

M.C.C.

CORTINARIUS (*HYDROCYBE*) GERMANUS. Fries.
under beech. Epping Forest, 1883.






CORTINARIUS (HYDROCYBE) OBTUSUS. Fries.
under trees, Monks Wood, Epping.


B



C.B.

CORTINARIUS (HYDROCYBE) ACUTUS. Fries.
in moist woods, near Bristol, &c.






CORTINARIUS (HYDROCYBE) JUNGHUHNII. Fries.
amongst moss. Northamptonshire.




strong scented

CORTINARIUS (HYDROCYBE) MILVINUS. Fries.
amongst leaves. Neasden, Oct. 1884.







CORTINARIUS (HYDROCYBE) DEPRESSUS. *Fries.*
in moist woods. *Cod. Cact., 1889.*






CORTINARIUS (*HYDROCYBE*) FASCIATUS. — Fries.
under fir trees. — Higwood Forest. — Oct. 1883.





GOMPHIDIUS GLUTINOSUS. Fries
under fir trees. Park End, Forest of Dean. Oct. 1857.




W.G.C.


18X7

W.G.S

GOMPHIDIUS ROSEUS Fries.
in woods. The Wrekin, Salop.









18-20 X 6


GOMPHIDIUS VISCIDUS Fries.
under fir trees. Credin Hill Camp. Sept. 1882.





H.G.B.


GOMPHIDIUS MACULATUS. Scop. (Variety)
near fir trees. Dinedor Camp.





G.M.

GOMPHIDIUS GRACILIS. Berk.
under fir trees. near Scarborough.





odour mealy, strong.



G.M.

PAXILLUS (LEPISTA) LEPISTA. Fries.
in fir woods. Scarborough.






PAXILLUS (LEPISTA) PANÆOLUS. Fries.
in mixed woods. Blaize Castle, Bristol.




G.M.


8X4

PAXILLUS (LEPISTA) ORCELLOIDES. C. G. M.
amongst grass. Queen's Cottage, Kew. Nov. 1886.






PAXILLUS (LEISTA) EXTENUATUS. Fries.
amongst grass. near Hereford. (Dr. Bull.)





PAXILLUS (LEPISTA) LIVIDUS. Cooke.
in woods. Leigh Down. Oct. 1882.




tomentose.


G.M.

PAXILLUS (LEPISTA) PARADOXUS. (Kulch.) Berk.
on the ground. Scarborough.




PAXILLUS (*TAPINULA*) INVOLUTUS. Fries.
on the ground. Kew Gardens. Oct. 1882.






PAXILLUS (*TAPINIA*) LEPTOPUS. Fries.
on stumps. Queen's Cottage, Kew.






PAXILLUS (TAPINIA) ATRO-TOMENTOSUS. Fries.
Pine woods near Woburn Station. Sept. 1872.



15-18 X 7-8 M


T.H.

PAXILLUS (TAPINIA) CRASSUS. Fries.
on mound of rifle butts, Blackheath. Nov. 1885.






PAXILLUS (*TAPINIA*) PANUOIDES. *Fries.*
on sawdust. *Coed Coch.*





HYGROPHORUS (*LIMACIUM*) CHRYSODON Fries.
under trees, Westridge Wood, near Nibley, Nov. 1880




mild.

2




HYGROPHORUS (*LIMACIUM*) EBURNEUS. Fries
in mixed woods. Epping Forest.



strong scented.

M.C.C.

HYGROPHORUS (*LIMACIUM*) COSSUS Fries.
in mixed woods. Dinnmore. Oct. 1887.




A




HYGROPHORUS PULVERULENTUS. B. & Br.
amongst pine leaves. Durdham Down.

B




7-8X4

HYGROPHORUS (LIMACIUM) PENARIUS. Fris.
in mixed woods. Morpeth, Oct. 1887.






HYGROPHORUS (*LIMACIUM*) ERUBESCENS Fries
in fir woods. near Hereford.




HYGROPHORUS (*Lutaceum*) PUDORINUS. — *Fries.*

in woods. — Crowell, Oxon, Oct. 1856.






HYGROPHORUS (*LIMACIUM*) GLUTINIFER Bull.
in mixed woods near Slough






HYGROPHORUS (LIMACIUM) ARBUSTIVUS. Fries.
under birches. Leigh Wood, Bristol.



HYGROPHORUS (LIMACIUM) AUREUS. Arrh.
amongst grass. Kew Gardens. Nov. 1883.






8X4

HYGROPHORUS (*LIMACIUM*) DISCOIDEUS. — Fries.
by roadsides in woods. — New Forest.





HYGROPHORUS (*LIMACIUM*) *LIMACINUS*. Fries.
in woods. Blaize Castle Woods, Nov. 1886.






HYGROPHORUS (*LIMACIUM*) OLIVACEO-ALBUS Fries
near fir trees. Tunel Wood, Watford. Oct. 1883.





HYGROPHORUS (LIMACIUM) HYPOTHEJUS Fries.
on grassy bank. near Dorking.



odour of cherry laurel.


margin flocculose.



W.G.S.

M.J.B.


HYGROPHORUS (*LIMACIUM*) CERASINUS. Berk.
amongst grass. Street, Somerset, 1868.





W.G.S.

HYGROPHORUS (*LIMACIUM*) FUSCO-ALBUS. — Lasch.
in mixed woods. — Jedburgh.



8
odour pleasant.



10X5

HYGROPHORUS (*LIMACIUM*) AGATHOSMUS. Fries.
in woods. Carlisle, Oct. 1886.






HYGROPHORUS (LIMACIUM) MESOTEFHRUS B. & Br
in damp places. Lyndhurst, Hants.






HYGROPHORUS (LIMACIUM) LIVIDO-ALBUS. Fries.
in grassy places. Queen's Cottage, Kew, Nov. 1886.



10 X 8 M

C.E.B.

HYGROPHORUS (CAMAROPHYLLUS) CAPRINUS. Scop.
under fir trees. near Bath.





W.G.S.

HYGROPHORUS (CAMAROPHYLLUS) LEPORINUS. Fries.
in woods. Forres, N.B.






W.G.S.


HYGROPHORUS (*CAMAROPHYLLUS*) NEMOREUS. Fries (?).


in pastures. Bishops Stortford, Oct. 1867.






HYGROPHORUS (CAMAROPHYLLUS) PRATENSIS. Pers.
amongst grass. Pleasure Grounds, Kew.





HYGROPHORUS (CAMAROPHYLLUS) PRATENSIS var. PALLIDUS. Berk.
in pastures. near Slough.

B



6X4


HYGROPHORUS (CAMAROPHYLLUS) PRATENSIS var. CINEREUS. Fries.
in pastures. near Slough, 1882.





HYGROPHORUS (*CAMAROPHYLLUS*) VIRGINEUS.
in grassy places. Cuckoo Pits, Epping Forest.

W. J. Wulf.
Oct. 1883.




gills thick.


G.M.

HYGROPHORUS (CAMAROPHYLLUS) VIRGINEUS, var. ROSEIPES
in fir woods, near Carlisle, Oct. 1887.






HYGROPHORUS (CAMAROPHYLLUS) NIVEUS. Scop.
amongst moss. Breinton, Sept. 1884.



R.E.B.

HYGROPHORUS (CAMAROPHYLLUS) RUSSO-CORIACEUS. B. & Br.
amongst grass. Ludlow, Coal Cock, &c.





Mrs. Russell

Kenilworth.

HYGROPHORUS (CAMAROPHYLLUS) VENTRICOSUS. B. & Br.

amongst grass. Coed Coch, and Kenilworth.


"*Hypomyces olivalis* sp. ?
+ *H. formicarum*"




W.G.S.

HYGROPHORUS (CAMAROPHYLLUS) FORNICATUS. Fries
amongst grass. Holm Lacey, Oct. 1872.



sweet scented like almond.


HYGROPHORUS (CAMAROPHYLLUS) DISTANS. — B. & Br.
in woods. — Addington Plantations, Oct. 1870.





HYGROPHORUS (CAMAROPHYLLUS) CLARKII. B. & Br.
amongst grass. Pleasure Grounds, Kew.

B



C.B.

HYGROPHORUS (CAMAROPHYLLUS) OVINUS. Bull.
in pastures, near Bristol.



mealy odour.




HYGROPHORUS MELEAGRINUS METAPODUS. — Pers.

n. sp. — F. 1907 — L. 1908 — S. 1908 —





HYGROPHORUS (CAMAROPHYLLUS) SUBRADIATUS. Fries.
in pastures, Cumberland.




HYGROPHORUS SUBRADIATUS var. LACMUS. Fries.
on healthy ground. near Hereford.




glutinous.


HYGROPHORUS (CAMAROPHYLLUS) IRRIGATUS. Fries.

amongst grass, after much rain. Kew Gardens, Oct. 1882.





HYGROPHORUS (HYGROCYBE) COLEMANNIANUS. Blox.
in field. Severn House, Henbury, Nov. 1877.




W.P.

HYGROPHORUS (HYGROCYBE) FETENS. Phil.
amongst grass. Shrewsbury





HYGROPHORUS (HYGROCYBE) SCIOPHANUS.
amongst moss. near Kendal. Fries.



HYGROPHORUS (HYGROCYBE) MUCRONELLUS. Fries.
in fields. near King's Lynn.

"Hyp. lactea pl 938 & H.
Houghtiana pl 936 seem
be the same species"



HYGROPHORUS (HYGROCYBE) LAETUS. Fries.
amongst grass. Epping Forest.






H.G.B.

HYGROPHORUS (CAMAROPHYLLUS) HOUGHTONI. B. & Br.
amongst grass. near Wellington, Salop.





A

HYGROPHORUS (HYGROCYBE) VITELLINUS.
amongst grass. Cassiobury Park, Watford.

Fries.
Nov. 1882.



B

W.G.S.

8 X 6

C.B.


HYGROPHORUS (HYGROCYBE) CEPACEUS Fries.
in grassy places. Coed Coch.



10-12 X 6 M

HYGROPHORUS (HYDROCYBE) COCCINEUS. Schaff.
in grassy places. Queen's Cottage, Kew.





HYGROPHORUS (HYGROCYBE) MINIATUS. Fries.
amongst grass. Kew Gardens, Nov. 1883.

B



HYGROPHORUS (HYGROCYBE) TURUNDUS. Fr. var. MOLLIS. Berk.
in grassy places. Coed Coch.



A

W.G.S.

7-8 X 6



M.J.B.

HYGROPHORUS (HYGROCYBE) WYNNIAE. *B. & Br.*
amongst twigs, &c. Coed Coch.


B

4 X 3



HYGROPHORUS (HYGROCYBE) MICACEUS. *B. & Br.*
on clay soil. Coed Coch.






8X5M


HYGROPHORUS (HYGROCYBE) PUNICEUS. Fries.
amongst moss, &c. Bedfordshire.





10-12 X 7-8


HYGROPHORUS (HYGROCYBE) OBRUSSEUS. Fries,
in grassy places. Oxford, Aug. 1886.



mealy odour.

HYGROPHORUS (*HYGROCYBE*) INTERMEDIUS Pass.


amongst grass, by realside, Langley, 1882.





10 X 7


HYGROPHORUS (HYGROCYBE) CONICUS. Fries.
on lawns. Kew Gardens.





M.C.C.

HYGROPHORUS (HYGROCYBE) CALYPTRÆFORMIS Bork.
amongst grass. Queen's Cottage, Kew.




fragile.



HYGROPHORUS (HYGROCYBE) CALYPTRÆFORMIS var. NIVEUS.


amongst grass. Queen's Cottage Grounds, Kew.






8 X 5


HYGROPHORUS (HYGROLYSI) CHLOROPHANUS. — Dr. S.
amongal grass. — Picture Grav'd, Kew, Oct. 1883.






HYGROPHORUS HYGROCYBE PSITTACINUS. Fr.
on lawns and pastures. Chingford, 1886.





HYGROPHORUS (HYGROCYBE) UNGUINOSUS. Fries.
in moist woods, near Hartford.



odour strong and disagreeable.



HYGROPHORUS (HYGROCYBE) NITRATUS. P.

amongst grass in woods. Ditcham Down, Oct 1882.



BINDING

MAR 18 1974

QK Cooke, Mordecai Cubitt
607 Illustrations of British
C67 Fungi
v.6

BioMed

PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET

UNIVERSITY OF TORONTO LIBRARY

